

CHEM 1423 - Exam 1 – February 11, 2016

Constants and Equations: $R = 8.31 \text{ J/mol-K}$

Beer-Lambert Law: $A = \log\left(\frac{I_o}{I}\right) = \epsilon bc$

Michaelis-Menten Equation: $v_0 = \frac{V_m[S]}{K_M + [S]}$

PERIODIC TABLE OF THE ELEMENTS

1 IA		2 IIA												16 VIA		17 VIIA		18 VIIIA																																																																																					
1	H 1.008	2	He 4.003	3	Li 6.941	4	Be 9.012	5	B 10.81	6	C 12.01	7	N 14.01	8	O 16.00	9	F 19.00	10	Ne 20.18	11	Na 23.00	12	Mg 24.30	13	Al 26.98	14	Si 28.09	15	P 30.97	16	S 32.07	17	Cl 35.45	18	Ar 39.95																																																																				
19	K 39.10	20	Ca 40.08	21	Sc 44.96	22	Ti 47.88	23	V 50.94	24	Cr 52.00	25	Mn 54.94	26	Fe 55.85	27	Co 58.93	28	Ni 58.69	29	Cu 63.55	30	Zn 65.38	31	Ga 69.72	32	Ge 72.59	33	As 74.92	34	Se 78.96	35	Br 79.90	36	Kr 83.80	37	Rb 85.47	38	Sr 87.62	39	Y 88.91	40	Zr 91.22	41	Nb 92.91	42	Mo 95.94	43	Tc (98)	44	Ru 101.1	45	Rh 102.9	46	Pd 106.4	47	Ag 107.9	48	Cd 112.4	49	In 114.8	50	Sn 118.7	51	Sb 121.8	52	Te 127.6	53	I 126.9	54	Xe 131.3	55	Cs 132.9	56	Ba 137.3	57	La 138.9	58	Ce 140.1	59	Pr 140.9	60	Nd 144.2	61	Pm (145)	62	Sm 150.4	63	Eu 152.0	64	Gd 157.2	65	Tb 158.9	66	Dy 162.5	67	Ho 164.9	68	Er 167.3	69	Tm 168.9	70	Yb 173.0
87	Fr (223)	88	Ra 226.0	89	Ac 227.0	90	Th 232.0	91	Pa 231.0	92	U 238.0	93	Np 237.0	94	Pu 244.0	95	Am 243.0	96	Cm 247.0	97	Bk 247.0	98	Cf 251.0	99	Es 252.0	100	Fm 257.0	101	Md 258.0	102	No 259.0	103	Lr 260.0	104	Rf 261.0	105	Db 262.0	106	Sg 263.0	107	Bh 262.0	108	Hs 265.0	109	Mt 266.0	110	Uu (267)	111	Uub (268)	112	Uuq (269)	113	Uuq (270)	114	Uuq (271)	115	Uuq (272)	116	Uuq (273)	117	Uuq (274)	118	Uuq (275)	119	Uuq (276)	120	Uuq (277)																																				

Lanthanide series											
57	La	58	Ce	59	Pr	60	Nd	61	Pm	62	Sm
89	Ac	90	Th	91	Pa	92	U	93	Np	94	Pu
Actinide series											

Key	
1	Atomic Number
H	Symbol
1.008	Atomic Mass

Approved by the I.U.P.A.C. Council in Geneva, August 30, 1997

Copyright © September • 1997 Educational Innovations, Inc.

CHEM 1423 - Exam 1 – February 11, 2016

Name _____

(60) PART I. MULTIPLE CHOICE (Circle the ONE correct answer)

1. Consider the hypothetical reaction, $2A + B \rightarrow 3C$. If the rate of change of $[C]$ is $\Delta[C]/dt = +0.60 \text{ M hr}^{-1}$. What is the rate of change of $[A]$, $\Delta[A]/dt$?
(A) -0.20 M hr^{-1} (B) -0.40 M hr^{-1}
(C) -0.30 M hr^{-1} (D) $+0.40 \text{ M hr}^{-1}$
2. The rate of the chemical reaction involving two substances, A and B, is measured. It is found that if the initial concentration of A used is tripled, keeping the B concentration the same, the rate increases by a factor of 27 (relative to the first experiment). If the concentrations of both A and B are doubled, the rate increases by a factor of 32 (relative to the first experiment). The rate law for this reaction is: Rate =
(A) $k[A]^2[B]^3$ (B) $k[A]^2[B]^2$ (C) $k[A]^3[B]$ (D) $k[A]^3[B]^2$

For #3 - #4: Consider a reaction, $A \rightarrow \text{Products}$, which is of order "n"; i.e. $\text{Rate} = k[A]^n$. For this reaction, the following initial rate data was obtained.

When $[A]_0 = 0.20 \text{ M}$, the initial rate is 0.50 M/s


When $[A]_0 = 0.60 \text{ M}$, the initial rate is 13.5 M/s

3. The order of this reaction (i.e. "n") is:
(A) +3 (B) +2 (C) +1 (D) -1
4. The rate constant for this reaction (i.e. "k") is:
(A) $313 \text{ M}^{-3}\text{s}^{-1}$ (B) $12.5 \text{ M}^{-1}\text{s}^{-1}$ (C) $62.5 \text{ M}^{-2}\text{s}^{-1}$ (D) 2.5 s^{-1}
5. Radioactive decay follows first order kinetics. Thallium-201 is a radioisotope which is used for myocardial imaging tests. Its half-life is 73 hours. If 300 mg of ^{201}Tl is injected into a patient, how many mg will remain 185 hours after the injection ?
(A) 124 mg (B) 52 mg (C) 34 mg (D) 83 mg.
6. For the **zeroth** order reaction, $A \rightarrow \text{Products}$, when the initial concentration of A is 0.90 M, it takes 25 s for the concentration to decrease to 0.30 M. Therefore, the rate constant for this reaction is approximately:
(A) 0.044 Ms^{-1} (B) 0.089 Ms^{-1} (C) 0.015 Ms^{-1} (D) 0.024 Ms^{-1}

For #7 - #8: Consider a reaction, $A \rightarrow \text{Products}$, which is of **second** order; i.e. $\text{Rate} = k[A]^2$. For this reaction, the rate constant is $8.5 \times 10^{-3} \text{ M}^{-1}\text{s}^{-1}$.

7. For this reaction, a plot of _____ vs. time is a straight line with a _____ slope.
- (A) $\ln([A]_t)$, negative (B) $[A]_t$, negative
(C) $1/[A]_t$, negative (D) $1/[A]_t$, positive
8. If the initial concentration of A is 0.75 M, what will be the approximate concentration of A 600 seconds after the start of the reaction.
- (A) 0.20 M (B) 0.16 M (C) 0.28 M (D) 0.046 M
9. Which **one** statement below is correct? The frequency factor in the Arrhenius equation can best be described as:
- (A) A numerical description of the amount of energy released by colliding reactant molecules when they form products.
- (B) The probability that a collision between molecules will have the correct relative orientation for reaction to occur.
- (C) A numerical description of how often molecules collide with the proper orientation to react at a specific concentration.
- (D) A numerical description of the amount of energy needed by colliding reactant molecules in order to form products.
10. Which of the following statements is/are **NOT** true?
- (1) The mechanism for a catalyzed reaction is different from the mechanism of the same reaction without the catalyst.
- (2) The enthalpy change of the reaction is lower for the catalyzed reaction than for the uncatalyzed chemical reaction.
- (3) The catalyst in a reaction is present at the start of the reaction and is regenerated during the reaction.
- (4) The Rate Determining Step in a catalyzed reaction mechanism has a higher activation energy and, therefore, is faster than the Rate Determining Step for the uncatalyzed reaction.
- (A) 2 & 4 (B) 2 & 3 & 4 (C) 1 & 3 (D) 2 & 4

11. For the reaction, $\text{Cd}_2^{2+}(\text{aq}) + \text{Zr}^{3+}(\text{aq}) \rightarrow 2 \text{Cd}^{2+}(\text{aq}) + \text{Zl}^+(\text{aq})$, the accepted reaction mechanism is:



An acceptable rate law for the rate, $\text{Rate} = \Delta[\text{Zr}^+]/\Delta t$, is:

- (A) $\text{Rate} = k' \frac{[\text{Zr}^{3+}][\text{Cd}_2^{2+}]}{[\text{Cd}]}$ (B) $\text{Rate} = k' \frac{[\text{Zr}^{3+}][\text{Cd}^+]^2}{[\text{Cd}^{2+}]}$
 (C) $\text{Rate} = k' [\text{Zr}^{3+}][\text{Cd}_2^{2+}][\text{Cd}^{2+}]$ (D) $\text{Rate} = k' \frac{[\text{Zr}^{3+}][\text{Cd}_2^{2+}]}{[\text{Cd}^{2+}]}$

12. Consider a reaction, $\text{R} \rightarrow \text{P}$ (i.e. Reactants \rightarrow Products). If the activation energy for the **reverse** reaction is 85 kJ/mol and the overall energy (aka enthalpy) change for the reaction is -30 kJ/mol. Therefore, the activation energy for the **forward** reaction is:

- (A) +115 kJ/mol (B) +55 kJ/mol (C) +105 kJ/mol (D) -55 kJ/mol

13. In an enzyme catalyzed reaction, when the Substrate Concentration is much greater than the Michaelis-Constant, K_M , then the rate of the reaction is _____ order with respect to $[\text{S}]$

- (A) Second order (B) First Order (C) Zeroth Order
 (D) Indeterminate without know the values of K_M and $[\text{S}]$.

14. In an enzyme catalyzed reaction, for what ratio, $[\text{S}]/K_M$, is the initial velocity, $v_0 = 0.70 V_m$?

- (A) 1.75 (B) 0.43 (C) 2.33 (D) 3.41

15. Consider the gas phase equilibrium, $\text{A}(\text{g}) \rightleftharpoons \text{B}(\text{g}) + 3\text{C}(\text{g})$,

$K_c = 1.0 \times 10^{-5}$. 3.0 mol of $\text{A}(\text{g})$ is placed in a 2.0 L container and the mixture is allowed to come to equilibrium. Calculate the approximate concentration of $\text{C}(\text{g})$ at equilibrium.

NOTE: You can assume that very little $\text{A}(\text{g})$ reacts to form $\text{B}(\text{g})$ and $\text{C}(\text{g})$

- (A) $8.2 \times 10^{-2} \text{ M}$ (B) $1.1 \times 10^{-1} \text{ M}$ (C) $2.7 \times 10^{-2} \text{ M}$ (D) $9.7 \times 10^{-2} \text{ M}$

For #16 - #20 For the gas phase reaction, $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{SO}_3(\text{g})$, $K_c = 120$. at 600 K. The Enthalpy change for this reaction is +95 kJ/mol

16. If a mixture is prepared with $[\text{SO}_2] = 0.4 \text{ M}$, $[\text{O}_2] = 0.3 \text{ M}$ and $[\text{SO}_3] = 1.5 \text{ M}$, the reaction quotient is _____ and the reaction will proceed towards the _____.


- (A) 47. , Right (B) 188. , Left
 (C) 63. , Right (D) 188. , Right

17. For the above reaction, if the equilibrium concentrations (at 600 K) of SO_3 and O_2 are 0.40 M and 0.10 M, respectively, what is the approximate equilibrium concentration of SO_2 ?
- (A) 0.12 M (B) 0.18 M (C) 0.013 M (D) 0.26 M
18. For the above equilibrium reaction, if the temperature is **decreased**, the ratio $[\text{SO}_2]/[\text{SO}_3]$ will _____ and K_c will _____.
- (A) decrease , increase (B) decrease , decrease
(C) increase , remain constant (D) increase , decrease
19. For the above reaction, if $\text{N}_2(\text{g})$ is added to the mixture in a container at fixed **volume**, the ratio $[\text{SO}_2]/[\text{SO}_3]$ will _____ and K_c will _____.
- (A) increase , decrease (B) decrease , remain constant
(C) remain constant , remain constant (D) increase , remain constant
20. What is the approximate equilibrium constant (at 600 K) for the related equilibrium, $4\text{SO}_2(\text{g}) + 2\text{O}_2(\text{g}) \rightleftharpoons 4\text{SO}_3(\text{g})$?
- (A) 240 (B) 1.4×10^4 (C) 6.9×10^{-5} (D) 8.3×10^{-3}
-

PART II. THERE ARE FOUR (4) PROBLEMS BELOW.

- (10) 1. The measured value of the rate constant for a first order reaction is $1.50 \times 10^3 \text{ min}^{-1}$ at a temperature of $120 \text{ }^\circ\text{C}$, and $1.35 \times 10^5 \text{ min}^{-1}$ at $180 \text{ }^\circ\text{C}$. Calculate the Arrhenius **Activation Energy** (in kJ/mol) and the **Frequency Factor** (in min^{-1}) for this reaction.

(10) 2. Consider a molecule, B, which reacts to form Products via First Order kinetics:



The molecule, B, absorbs visible light at 480 nm. The Molar Absorptivity for this absorption is given by $\varepsilon = 800 \text{ M}^{-1} \text{ cm}^{-1}$.

When a sample of B is prepared in a 0.5 cm cell, the Percent Transmission at the beginning of the reaction was 25%. After the reaction has run for 50 minutes, the percent transmission is 70%. Calculate the rate constant for this reaction (**give the units**)

(10) 3. Consider the reaction: $\text{Cl}_2(\text{g}) + 2 \text{NO}_2(\text{g}) \rightleftharpoons 2 \text{NOCl}(\text{g})$.

If 1.50 mol of $\text{Cl}_2(\text{g})$ and 1.50 mol of $\text{NO}_2(\text{g})$ are placed in a 5.0 Liter container, which is heated to $200\text{ }^\circ\text{C}$, the equilibrium concentration of $\text{NO}_2(\text{g})$ is 0.120 M. Calculate the equilibrium constant, K_c , for this reaction at $200\text{ }^\circ\text{C}$.

- (10) 4. Consider the reaction: $H_2(g) + I_2(g) \xrightleftharpoons{K_c} 2HI(g)$ $K_c = 9.0$ (at 550 K)
8.0 mol of HI(g) is placed in a 20.0 L container and heated to 550 K, where equilibrium is established.

Calculate the equilibrium concentrations (**in M**) of $H_2(g)$, $I_2(g)$ and $HI(g)$?