

# CHEM 1423 - Exam 1 – February 9, 2017

Constants and Equations:  $R = 8.31 \text{ J/mol-K}$

Beer-Lambert Law:  $A = \log\left(\frac{I_o}{I}\right) = \epsilon bc$

Michaelis-Menten Equation:  $v_0 = \frac{V_m[S]}{K_M + [S]}$

## PERIODIC TABLE OF THE ELEMENTS

1 IA		2 IIA												16 VIA		17 VIIA		18 VIIIA																																																																																					
1	<b>H</b> 1.008	2	<b>He</b> 4.003	3	<b>Li</b> 6.941	4	<b>Be</b> 9.012	5	<b>B</b> 10.81	6	<b>C</b> 12.01	7	<b>N</b> 14.01	8	<b>O</b> 16.00	9	<b>F</b> 19.00	10	<b>Ne</b> 20.18	11	<b>Na</b> 23.00	12	<b>Mg</b> 24.30	13	<b>Al</b> 26.98	14	<b>Si</b> 28.09	15	<b>P</b> 30.97	16	<b>S</b> 32.07	17	<b>Cl</b> 35.45	18	<b>Ar</b> 39.95																																																																				
19	<b>K</b> 39.10	20	<b>Ca</b> 40.08	21	<b>Sc</b> 44.96	22	<b>Ti</b> 47.88	23	<b>V</b> 50.94	24	<b>Cr</b> 52.00	25	<b>Mn</b> 54.94	26	<b>Fe</b> 55.85	27	<b>Co</b> 58.93	28	<b>Ni</b> 58.69	29	<b>Cu</b> 63.55	30	<b>Zn</b> 65.38	31	<b>Ga</b> 69.72	32	<b>Ge</b> 72.59	33	<b>As</b> 74.92	34	<b>Se</b> 78.96	35	<b>Br</b> 79.90	36	<b>Kr</b> 83.80	37	<b>Rb</b> 85.47	38	<b>Sr</b> 87.62	39	<b>Y</b> 88.91	40	<b>Zr</b> 91.22	41	<b>Nb</b> 92.91	42	<b>Mo</b> 95.94	43	<b>Tc</b> (98)	44	<b>Ru</b> 101.1	45	<b>Rh</b> 102.9	46	<b>Pd</b> 106.4	47	<b>Ag</b> 107.9	48	<b>Cd</b> 112.4	49	<b>In</b> 114.8	50	<b>Sn</b> 118.7	51	<b>Sb</b> 121.8	52	<b>Te</b> 127.6	53	<b>I</b> 126.9	54	<b>Xe</b> 131.3	55	<b>Cs</b> 132.9	56	<b>Ba</b> 137.3	57	<b>La</b> 138.9	58	<b>Ce</b> 140.1	59	<b>Pr</b> 140.9	60	<b>Nd</b> 144.2	61	<b>Pm</b> (145)	62	<b>Sm</b> 150.4	63	<b>Eu</b> 152.0	64	<b>Gd</b> 157.2	65	<b>Tb</b> 158.9	66	<b>Dy</b> 162.5	67	<b>Ho</b> 164.9	68	<b>Er</b> 167.3	69	<b>Tm</b> 168.9	70	<b>Yb</b> 173.0
7	<b>Fr</b> (223)	8	<b>Ra</b> 226.0	9	<b>Ac</b> 227.0	10	<b>Th</b> 232.0	11	<b>Pa</b> 231.0	12	<b>U</b> 238.0	13	<b>Np</b> 237.0	14	<b>Pu</b> 244.0	15	<b>Am</b> 243.0	16	<b>Cm</b> 247.0	17	<b>Bk</b> 247.0	18	<b>Cf</b> 251.0	19	<b>Es</b> 252.0	20	<b>Fm</b> 257.0	21	<b>Md</b> 258.0	22	<b>No</b> 259.0	23	<b>Lr</b> 260.0	24	<b>Rf</b> 261.0	25	<b>Db</b> 262.0	26	<b>Sg</b> 263.0	27	<b>Bh</b> 262.0	28	<b>Hs</b> 265.0	29	<b>Mt</b> 266.0	30	<b>Hf</b> 178.5	31	<b>Ta</b> 180.9	32	<b>W</b> 183.8	33	<b>Re</b> 186.2	34	<b>Os</b> 190.2	35	<b>Ir</b> 192.2	36	<b>Pt</b> 195.1	37	<b>Au</b> 197.0	38	<b>Hg</b> 200.6	39	<b>Tl</b> 204.4	40	<b>Pb</b> 207.2	41	<b>Bi</b> 209.0	42	<b>Po</b> (209)	43	<b>At</b> (210)	44	<b>Rn</b> (222)																												

Lanthanide series																											
57	<b>La</b>	58	<b>Ce</b>	59	<b>Pr</b>	60	<b>Nd</b>	61	<b>Pm</b>	62	<b>Sm</b>	63	<b>Eu</b>	64	<b>Gd</b>	65	<b>Tb</b>	66	<b>Dy</b>	67	<b>Ho</b>	68	<b>Er</b>	69	<b>Tm</b>	70	<b>Yb</b>
138.9	140.1	140.9	144.2	(145)	150.4	152.0	157.2	158.9	162.5	164.9	167.3	168.9	173.0	173.0	173.0	173.0	173.0	173.0	173.0	173.0	173.0	173.0	173.0	173.0	173.0	173.0	173.0
Actinide series																											
89	<b>Ac</b>	90	<b>Th</b>	91	<b>Pa</b>	92	<b>U</b>	93	<b>Np</b>	94	<b>Pu</b>	95	<b>Am</b>	96	<b>Cm</b>	97	<b>Bk</b>	98	<b>Cf</b>	99	<b>Es</b>	100	<b>Fm</b>	101	<b>Md</b>	102	<b>No</b>
227.0	232.0	231.0	238.0	237.0	244.0	243.0	247.0	247.0	251.0	252.0	257.0	258.0	259.0	259.0	259.0	259.0	259.0	259.0	259.0	259.0	259.0	259.0	259.0	259.0	259.0		

Approved by the I.U.P.A.C. Council in Geneva, August 30, 1997

Copyright © September, 1997 Educational Innovations, Inc.

CHEM 1423 - Exam 1 – February 9, 2017

Name \_\_\_\_\_

(76) PART I. MULTIPLE CHOICE (Circle the ONE correct answer)

1. For the reaction,  $A + B \rightarrow \text{Products}$ , the rate law is  $\text{Rate} = k \frac{[A]}{[B]^2}$ . The units of the rate constant are:
- (A)  $M^3s^{-1}$                       (B)  $M^2s^{-1}$                       (C)  $M^{-3}s^{-1}$                       (D)  $M^{-2}s^{-1}$
2. The rate of the chemical reaction involving two substances, A and B, is measured. It is found that if the initial concentration of A used is quadrupled, keeping the B concentration the same, the rate increases by a factor of 64 (relative to the first experiment). If the concentrations of both A and B are quadrupled, the rate increases by a factor of 16 (relative to the first experiment). The rate law for this reaction is: Rate =
- (A)  $k[A]^2[B]$                       (B)  $k[A]^2[B]^{-1}$                       (C)  $k[A]^3[B]^{-1}$                       (D)  $k[A]^3[B]$
3. Consider a reaction,  $A \rightarrow \text{Products}$ , which is of order "n"; i.e.  $\text{Rate} = k[A]^n$ . For this reaction, the following initial rate data was obtained.
- When  $[A]_0 = 0.20 \text{ M}$ , the initial rate is  $1.2 \text{ M/s}$   
When  $[A]_0 = 0.80 \text{ M}$ , the initial rate is  $19.2 \text{ M/s}$
- The order of this reaction (i.e. "n") is:
- (A) +3                      (B) +2                      (C) +1                      (D) -1
4. For the above reaction (question immediately above), the rate constant is approximately:
- (A)  $6.0 \text{ s}^{-1}$                       (B)  $150 \text{ M}^{-2}\text{s}^{-1}$                       (C)  $30 \text{ M}^{-1}\text{s}^{-1}$                       (D)  $0.24 \text{ M}^2\text{s}^{-1}$

**For #5 - #7:** Consider a reaction,  $A \rightarrow \text{Products}$ , which is of **first** order; i.e.  $\text{Rate} = k[A]$ . For this reaction, the rate constant is  $0.015 \text{ s}^{-1}$  at  $100 \text{ }^\circ\text{C}$ . The Activation Energy for this reaction is  $75 \text{ kJ/mol}$ .

5. For this reaction, a plot of \_\_\_\_\_ vs. time is a straight line with a \_\_\_\_\_ slope.
- (A)  $\ln([A]_t)$ , negative                      (B)  $[A]_t$ , negative  
(C)  $1/[A]_t$ , negative                      (D)  $1/[A]_t$ , positive

6. If the initial concentration of A is 1.30 M (at 100 °C), what will be the concentration of A 70 s after the start of the reaction?  
(A) 0.25 M                      (B) 0.36 M                      (C) 0.45 M                      (D) 0.58 M
7. What will be the value of the rate constant at 150 °C?  
(A) 0.26 s<sup>-1</sup>    (B) 0.057 s<sup>-1</sup>    (C) 8.6x10<sup>-4</sup> s<sup>-1</sup>    (D) 17.5 s<sup>-1</sup>

**For #8 - #9:** Consider a reaction, B → Products, which is 2nd. order; i.e. Rate = k[B]<sup>2</sup>. The molecule, B, absorbs visible light at 500 nm. The Molar Absorptivity for this absorption is  $\epsilon = 600 \text{ M}^{-1} \text{ cm}^{-1}$ .



When a sample of B with initial concentration, [B]<sub>0</sub> = 1.20x10<sup>-3</sup> M, is placed in a sample cell with cell pathlength = 2.0 cm, then the Percent Transmission 150 s after the start of the experiment is 30%.

8. The concentration of [B] 150 s after the start of the experiment is approximately:  
(A) 4.36x10<sup>-4</sup> M    (A) 1.82x10<sup>-4</sup> M    (A) 1.00x10<sup>-3</sup> M    (A) 8.71x10<sup>-4</sup> M
9. The rate constant for this second order reaction is approximately:  
(A) 1.1 M<sup>-1</sup>s<sup>-1</sup>    (B) 2.1 M<sup>-1</sup>s<sup>-1</sup>    (C) 31.1 M<sup>-1</sup>s<sup>-1</sup>    (D) 9.7 M<sup>-1</sup>s<sup>-1</sup>
10. Which of the following statements is/are **TRUE**?
- (1) The mechanism for a catalyzed reaction is the same as the mechanism of the same reaction without the catalyst.
  - (2) The enthalpy change of the reaction is the same for the catalyzed reaction as for the uncatalyzed chemical reaction.
  - (3) The intermediate in a reaction is generated in one of the earlier steps of a reaction and used up in later steps.
  - (4) The Rate Determining Step in a catalyzed reaction mechanism has a lower activation energy and, therefore, is slower than the Rate Determining Step for the uncatalyzed reaction.
- (A) 1 & 4                      (B) 2 only                      (C) 2 & 3                      (D) 2 & 3 & 4

11. Consider a reaction,  $R \rightarrow P$  (i.e. Reactants  $\rightarrow$  Products). If the activation energy for the **forward** reaction is 40 kJ/mol and the activation energy for the **reverse** reaction is 85 kJ/mol, then the overall enthalpy (aka energy) change for this reaction is:

(A) +125 kJ/mol      (B) -45 kJ/mol      (C) -125 kJ/mol      (D) +45 kJ/mol

12. For the reaction,  $\text{Sc}^{3+}(\text{aq}) + 2 \text{Pd}^+(\text{aq}) \rightarrow \text{Sc}^+(\text{aq}) + 2 \text{Pd}^{2+}(\text{aq})$ , the reaction mechanism is:



The overall rate equation for this reaction is:

- (A)  $\text{Rate} = k' \frac{[\text{Pd}^{2+}][\text{Sc}^{3+}]}{[\text{Pd}^+]^2}$       (B)  $\text{Rate} = k' \frac{[\text{Pd}^+][\text{Sc}^{3+}]}{[\text{Pd}^{2+}]}$   
 (C)  $\text{Rate} = k' \frac{[\text{Pd}^+]^2[\text{Sc}^{3+}]}{[\text{Pd}^{2+}]}$       (D)  $\text{Rate} = k' [\text{Pd}][\text{Sc}^{3+}]$

13. When a substrate (S) binds **Strongly** to an enzyme (E) to form the complex, ES:

(A)  $K_m$  is small      (B)  $V_m$  is small      (C)  $V_m$  is large      (D)  $K_m$  is large

14. In an enzyme catalyzed reaction, for approximately what ratio,  $[\text{S}]/K_m$ , does one find that  $v_o = 0.4V_m$  ?

(A)  $[\text{S}]/K_m = 1.50$       (B)  $[\text{S}]/K_m = 1.20$       (C)  $[\text{S}]/K_m = 0.83$       (D)  $[\text{S}]/K_m = 0.67$

15. Consider the gas phase equilibrium,  $\text{A}(\text{g}) \rightleftharpoons \text{B}(\text{g}) + 2\text{C}(\text{g})$ ,

$K_c = 1.0 \times 10^{-4}$ . 3.0 mol of  $\text{A}(\text{g})$  is placed in a 4.0 L container and the mixture is allowed to come to equilibrium. Calculate the approximate concentration of  $\text{C}(\text{g})$  at equilibrium.

**NOTE: You can assume that very little  $\text{A}(\text{g})$  reacts to form  $\text{B}(\text{g})$  and  $\text{C}(\text{g})$**

(A)  $2.7 \times 10^{-2} \text{ M}$       (B)  $5.3 \times 10^{-2} \text{ M}$       (C)  $4.2 \times 10^{-2} \text{ M}$       (D)  $8.4 \times 10^{-2} \text{ M}$

16. Consider the gas phase equilibrium  $A(g) \rightleftharpoons 2 B(g)$ . 3.0 moles of pure  $A(g)$  are placed in a 2.0 L container, and the reaction is allowed to proceed to equilibrium. It is found that after equilibrium has been established, the concentration of  $B(g)$  is 0.80 M. The value of the equilibrium constant,  $K_c$ , is approximately:

- (A) 0.15                      (B) 0.43                      (C) 0.58                      (D) 0.73

**For #17 - #19:** For the gas phase reaction,  $2 Br_2(g) + 4 NO(g) \rightleftharpoons 4 NOBr(g)$ ,  $K_c = 50$ . at 400 K.

17. For the above reaction, if the equilibrium concentrations (at 400 K) of  $Br_2(g)$  and  $NOBr(g)$  are each 2.5 M, then the equilibrium concentration of  $NO$  is approximately:

- (A) 1.7 M                      (B) 2.8 M                      (C) 0.59 M                      (D) 0.35

18. If a mixture is prepared with  $[Br_2] = 0.7 M$ ,  $[NO] = 0.7 M$  and  $[NOBr] = 1.5 M$ , the reaction quotient is approximately \_\_\_\_\_ and the reaction will proceed towards the \_\_\_\_\_.

- (A) 43. , Right                      (B) 84 , Left                      (C) 43 , Left  
(D) None of the above

19. The equilibrium constant for the related reaction  $2 NOBr(g) \rightleftharpoons Br_2(g) + 2 NO(g)$ , at 400 K is approximately:

- (A) 0.14                      (B) 0.32                      (C) 0.04                      (D) 7.1

---

**PART II. THERE ARE TWO (2) PROBLEMS ON FOLLOWING PAGES**  
**You MUST show your work for credit.**

- (12) 1. Consider the reaction,  $A \rightarrow \text{Products}$ , which is **third** order with respect to  $[A]$ ; i.e. the rate is given by  $\text{Rate} = k[A]^3$ . It can be shown that the integrated rate equation for this reaction is given by:

The integrated rate equation for the reaction is: 
$$\frac{1}{[A]^2} - \frac{1}{[A]_0^2} = 2kt$$

$[A]_0$  and  $[A]$  are the concentrations at  $t = 0$  and  $t$ , respectively, and  $k$  is the rate constant.

- (8) (a) When the initial concentration of  $A$  is  $0.90 \text{ M}$ , the half-life for the reaction is  $t_{1/2} = 60 \text{ s}$ . Calculate the rate constant for the reaction (give units).

- (4) (b) When the initial concentration of  $A$  is  $0.90 \text{ M}$ , calculate the concentration of  $A$   $100 \text{ s}$  after the start of the reaction.

**Note:** If you don't like your answer for part (a), you can use  $k = 0.035 \text{ M}^{-2}\text{s}^{-1}$  (without loss of credit in this part).

(12) 2. Consider the reaction:  $2NO(g) \xrightleftharpoons{K_c} N_2(g) + O_2(g)$ . The equilibrium constant is  $K_c = 2$ . at 1500 K.

2.0 mol of  $N_2(g)$  and 2.0 mol of  $O_2$  are placed in a 10 L container and heated to 1500 K, where equilibrium is established.

Calculate the equilibrium concentrations (**in M**) of NO,  $N_2$  and  $O_2$  in the equilibrium mixture.