

CHEM 1423 - Exam 2 – March 2, 2017 - **Version A**

Constants and Conversion Factors

$$R = 0.082 \text{ L-atm/mol-K}$$

$$R = 8.31 \text{ J/mol-K}$$

$$1 \text{ atm.} = 760 \text{ torr}$$

Molar Masses: $\text{C}_2\text{H}_6\text{O}_2$ - 62. H_2O - 18. $\text{C}_6\text{H}_{12}\text{O}_6$ - 180.
 NH_3 - 17, $\text{C}_6\text{H}_5\text{C}_2\text{H}_5(\text{l})$ - 106

Beer-Lambert Law: $A = \log\left(\frac{I_0}{I}\right) = \epsilon bc$

CHEM 1423 - Exam 2 – March 2, 2017 - Version A

Name _____

(76) PART I. MULTIPLE CHOICE (Circle the ONE correct answer)

For #1 - #3: Consider the gas phase reaction, $2 \text{Br}_2(\text{g}) + 4 \text{NO}(\text{g}) \rightleftharpoons 4 \text{NOBr}(\text{g})$, $K_c = 50$. at 400 K. The enthalpy change for this reaction is $\Delta H = +75 \text{ kJ}$

- For the above equilibrium reaction, if $\text{NO}(\text{g})$ is added to the mixture, the ratio $[\text{NOBr}]/[\text{Br}_2]$ will _____ and K_c will _____.
(A) decrease , remain constant (B) increase , decrease
(C) increase , remain constant (D) decrease , decrease
- For the above equilibrium reaction, if the temperature is **decreased**, the ratio $[\text{NOBr}]/[\text{Br}_2]$ will _____ and K_c will _____.
(A) decrease , remain constant (B) increase , decrease
(C) increase , remain constant (D) decrease , decrease
- For the above reaction, if $\text{Ar}(\text{g})$ is added to the mixture in a container at fixed **total pressure**, the ratio $[\text{NOBr}]/[\text{Br}_2]$ will _____ and K_c will _____.
(A) decrease , remain constant (B) increase , decrease
(C) remain constant , remain constant (D) increase , remain constant
- Consider the equilibrium, $\text{H}_2(\text{gas}) + \text{I}_2(\text{solid}) \rightleftharpoons 2 \text{HI}(\text{gas})$
If the volume of the container is **decreased**, the ratio, $[\text{HI}(\text{g})]/[\text{H}_2(\text{g})]$, will _____ and K_c will _____.
(A) decrease , decrease (B) decrease , remain constant
(C) remain constant , remain constant (D) increase , remain constant

For #5-#6: Consider the aqueous solution equilibrium, $\text{A}(\text{aq}) \rightleftharpoons 3 \text{B}(\text{aq})$.
The product, B, has an absorption in the UV range of the spectrum at 450 nm, with a Molar Absorptivity, $\epsilon = 50. \text{ M}^{-1} \text{ cm}^{-1}$

A solution is prepared in a 1.5 cm cell with an initial concentration of the reactant, A, $[\text{A}]_0 = 0.005 \text{ M}$, and the solution is allowed to reach equilibrium.
At equilibrium, the % transmittance of B is 30%.

- What is the approximate concentration of B at equilibrium?
(A) 0.021 M (B) 0.0023 M (C) 0.0070 M (D) 0.00023 M

Version A

6. What is the approximate value of the equilibrium constant for the above reaction?
(A) 1.3×10^{-4} (B) 2.6 (C) 1.2×10^{-8} (D) 3.7×10^{-2}
7. Which of the following statements is/are **NOT correct**.
(i) the solubility of most solids in a liquid increases with rising temperature.
(ii) the solubility of most gases in a liquid increases with rising temperature.
(iii) when a solid is dissolved in a liquid, the entropy increases.
(iv) ΔH_{soln} must be negative for a solid to dissolve in a liquid.
(A) ii only (B) iv only (C) i & iii (D) ii & iv
8. A sample of water contains of Arsenic in a sample of water is 16 ppb (parts per billion) of Arsenic. Therefore, the Weight Percent of Arsenic in the sample is:
(A) $1.6 \times 10^{-10} \%$ (B) $1.6 \times 10^{-6} \%$ (C) $1.6 \times 10^{-9} \%$ (D) $1.6 \times 10^{-8} \%$
- For #9 - #10:** When 124 grams of Ethylene Glycol ($\text{C}_2\text{H}_6\text{O}_2$), is added to 600 grams of water, the density of the solution is 0.80 g/mL.
9. The **Molarity** of Ethylene Glycol in the above solution is approximately:
(A) 3.3 M (B) 2.7 M (C) 2.2 M (D) 3.5 M
10. The **mole fraction** of Ethylene Glycol in the above solution is approximately:
(A) 0.21 (B) 0.057 (C) 0.060 (D) 0.17
11. You want to prepare a 1.5 **molal** solution of Ethylene Glycol ($\text{C}_2\text{H}_6\text{O}_2$) in water. Approximately how many grams of Ethylene Glycol would you have to add to 600 grams of water to prepare this solution?
(A) 56 g (B) 65 g (C) 80 g (D) 41 g
12. What is the approximate weight % of Glucose $\text{C}_6\text{H}_{12}\text{O}_6$ in an aqueous solution containing 0.80 **molal** Glucose?
(A) 0.14 % (B) 1.4 % (C) 14.4 % (D) 12.6%
13. What is the approximate NH_3 **Molarity** in a solution in which the NH_3 mass percent is 10% (solution density = 0.92 g/mL)?
(A) 6.5 M (B) 5.4 M (C) 3.7 M (D) 5.9 M

Version A

14. When 60 grams of an unknown compound is dissolved in 500 g of water ($K_f=1.9\text{ }^\circ\text{C/m}$), the freezing point of the solution is $-2.20\text{ }^\circ\text{C}$. The Molar Mass of the compound is approximately
- (A) 52 g/mol (B) 104 g/mol (C) 70 g/mol (D) 86 g/mol
15. What is the osmotic pressure, **in torr**, when 9.5×10^{-4} mol of the strong electrolyte, aluminum nitrate $[\text{Al}(\text{NO}_3)_3]$, is dissolved in 600 mL of aqueous solution at $25\text{ }^\circ\text{C}$?
- (A) 0.16 torr (B) 29 torr (C) 118 torr (D) 105 torr
16. When 2.0 grams of an Enzyme are dissolved in 600 mL of aqueous solution, the osmotic pressure at $25\text{ }^\circ\text{C}$ is 9.5 torr. The Molar Mass of the Enzyme is approximately:
- (A) 6.5×10^3 g/mol (B) 4.4×10^3 g/mol (C) 2.3×10^3 g/mol
(D) Cannot be determined without the Osmotic Pressure Depression Constant
17. The vapor pressure of pure water at $60\text{ }^\circ\text{C}$ is 149 torr. What is the approximate vapor pressure of a solution prepared by adding 135 grams of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) to 300 grams of water at $60\text{ }^\circ\text{C}$?
- (A) 137.4 torr (B) 102.8 torr (C) 6.4 torr (D) 142.6 torr
18. Consider the strong electrolytes, sodium sulfate, Na_2SO_4 , and sodium phosphate, Na_3PO_4 .
- If 350 mL of 0.30 M $\text{Na}_2\text{SO}_4(\text{aq})$ is added to 250 mL of 0.50 M $\text{Na}_3\text{PO}_4(\text{aq})$, the sodium ion concentration (i.e Molarity) in the mixture, $[\text{Na}^+]$, is approximately:
- (A) 0.59 M (B) 0.98 M (C) 0.46 M (D) 0.35 M
19. Which one of the following solutions has the **lowest** boiling point?
- (A) 0.10 m $\text{Mg}_3(\text{PO}_4)_2$ (B) 0.11 m K_3AsO_4
(C) 0.20 m Na_2SO_4 (D) 0.32 m NH_4Cl

PART II. TWO (2) PROBLEMS ON FOLLOWING PAGES:

REMEMBER TO SHOW YOUR WORK FOR CREDIT

Version A

- (12) 1. Consider the equilibrium between $\text{N}_2(\text{g})$, $\text{H}_2(\text{g})$, $\text{NH}_3(\text{g})$:
 $\text{N}_2(\text{g}) + 3 \text{H}_2(\text{g}) \rightleftharpoons 2 \text{NH}_3(\text{g})$. The value of K_c at 30°C is 8.0. The Enthalpy change for this reaction is $\Delta H = -92.0 \text{ kJ}$.

Calculate the temperature, **in $^\circ\text{C}$** , at which the equilibrium constant, K_c , is 1.00×10^{-4} .

- (12) 2. The vapor pressure of pure Ethylbenzene, $\text{C}_6\text{H}_5\text{C}_2\text{H}_5(\text{l})$ [$M=106$], is 74.0 torr at 70°C . When 50 grams of an unknown non-volatile solute, X, is added to 265 grams of Ethylbenzene, the vapor pressure of the solution at 70°C is 62.9 torr.

Calculate the Molar Mass of the unknown, X, in grams/mol

CHEM 1423 - Exam 2 – March 2, 2017 - **Version B**

Constants and Conversion Factors

$$R = 0.082 \text{ L-atm/mol-K}$$

$$R = 8.31 \text{ J/mol-K}$$

$$1 \text{ atm.} = 760 \text{ torr}$$

Molar Masses: $\text{C}_2\text{H}_6\text{O}_2$ - 62. H_2O - 18. $\text{C}_6\text{H}_{12}\text{O}_6$ - 180.
 NH_3 - 17, $\text{C}_6\text{H}_5\text{C}_2\text{H}_5(\text{l})$ - 106

Beer-Lambert Law: $A = \log\left(\frac{I_0}{I}\right) = \epsilon bc$

CHEM 1423 - Exam 2 – March 2, 2017 - **Version B**

Name _____

(76) **PART I. MULTIPLE CHOICE (Circle the ONE correct answer)**

For #1-#2: Consider the aqueous solution equilibrium, $A(aq) \rightleftharpoons 3 B(aq)$.
The product, B, has an absorption in the UV range of the spectrum at 450 nm, with a Molar Absorptivity, $\epsilon = 50. \text{ M}^{-1} \text{ cm}^{-1}$

A solution is prepared in a 1.5 cm cell with an initial concentration of the reactant, A, $[A]_0 = 0.005 \text{ M}$, and the solution is allowed to reach equilibrium.
At equilibrium, the % transmittance of B is 30%.

1. What is the approximate concentration of B at equilibrium?
(A) 0.0070 M (B) 0.0023 M (C) 0.021 M (D) 0.00023 M
2. What is the approximate value of the equilibrium constant for the above reaction?
(A) 1.2×10^{-8} (B) 2.6 (C) 1.3×10^{-4} (D) 3.7×10^{-2}

For #3 - #5: Consider the gas phase reaction, $2 \text{ Br}_2(g) + 4 \text{ NO}(g) \rightleftharpoons 4 \text{ NOBr}(g)$,
 $K_c = 50.$ at 400 K. The enthalpy change for this reaction is $\Delta H = +75 \text{ kJ}$

3. For the above equilibrium reaction, if the temperature is **decreased**, the ratio $[\text{NOBr}]/[\text{Br}_2]$ will _____ and K_c will _____.
(A) increase , remain constant (B) increase , decrease
(C) decrease , remain constant (D) decrease , decrease
4. For the above equilibrium reaction, if $\text{NO}(g)$ is added to the mixture, the ratio $[\text{NOBr}]/[\text{Br}_2]$ will _____ and K_c will _____.
(A) decrease , remain constant (B) decrease , decrease
(C) increase , remain constant (D) increase , decrease
5. For the above reaction, if $\text{Ar}(g)$ is added to the mixture in a container at fixed **total pressure**, the ratio $[\text{NOBr}]/[\text{Br}_2]$ will _____ and K_c will _____.
(A) increase , remain constant (B) increase , decrease
(C) remain constant , remain constant (D) decrease , remain constant

Version B

6. Consider the equilibrium, $\text{H}_2(\text{gas}) + \text{I}_2(\text{solid}) \rightleftharpoons 2 \text{HI}(\text{gas})$
If the volume of the container is **decreased**, the ratio, $[\text{HI}(\text{g})]/[\text{H}_2(\text{g})]$, will _____
and K_c will _____
- (A) decrease , decrease (B) increase , remain constant
(C) remain constant , remain constant (D) decrease , remain constant

7. Which of the following statements is/are **NOT correct**.
- (i) the solubility of most solids in a liquid increases with rising temperature.
(ii) the solubility of most gases in a liquid .increases with rising temperature.
(iii) when a solid is dissolved in a liquid, the entropy increases.
(iv) ΔH_{soln} must be negative for a solid to dissolve in a liquid.
- (A) ii & iv (B) iv only (C) i & iii (D) ii only

For #8 - #9: When 124 grams of Ethylene Glycol ($\text{C}_2\text{H}_6\text{O}_2$), is added to 600 grams of water, the density of the solution is 0.80 g/mL.

8. **Molarity** of Ethylene Glycol in the above solution is approximately:
- (A) 2.2 M (B) 2.7 M (C) 3.3 M (D) 3.5 M
9. The **mole fraction** of Ethylene Glycol in the above solution is approximately:
- (A) 0.21 (B) 0.057 (C) 0.060 (D) 0.057
10. A sample of water contains of Arsenic in a sample of water is 16 ppb (parts per billion) of Arsenic. Therefore, the Weight Percent of Arsenic in the sample is:
- (A) $1.6 \times 10^{-10} \%$ (B) $1.6 \times 10^{-9} \%$ (C) $1.6 \times 10^{-6} \%$ (D) $1.6 \times 10^{-8} \%$
11. You want to prepare a 1.5 **molal** solution of Ethylene Glycol ($\text{C}_2\text{H}_6\text{O}_2$) in water. Approximately how many grams of Ethylene Glycol would you have to add to 600 grams of water to prepare this solution?
- (A) 80 g (B) 65 g (C) 56 g (D) 41 g
12. What is the approximate NH_3 **Molarity** in a solution in which the NH_3 mass percent is 10% (solution density = 0.92 g/mL)?
- (A) 6.5 M (B) 5.9 M (C) 3.7 M (D) 5.4 M
13. What is the approximate weight % of Glucose $\text{C}_6\text{H}_{12}\text{O}_6$ in an aqueous solution containing 0.80 **molal** Glucose?
- (A) 12.6 % (B) 1.4 % (C) 14.4 % (D) 0.14%

Version B

14. When 2.0 grams of an Enzyme are dissolved in 600 mL of aqueous solution, the osmotic pressure at 25 °C is 9.5 torr. The Molar Mass of the Enzyme is approximately:
- (A) 4.4×10^3 g/mol (B) 6.5×10^3 g/mol (C) 2.3×10^3 g/mol
(D) Cannot be determined without the Osmotic Pressure Depression Constant
15. What is the osmotic pressure, **in torr**, when 9.5×10^{-4} mol of the strong electrolyte, aluminum nitrate $[\text{Al}(\text{NO}_3)_3]$, is dissolved in 600 mL of aqueous solution at 25 °C?
- (A) 0.16 torr (B) 29 torr (C) 105 torr (D) 118 torr
16. When 60 grams of an unknown compound is dissolved in 500 g of water ($K_f = 1.9$ °C/m), the freezing point of the solution is -2.20 °C. The Molar Mass of the compound is approximately
- (A) 104 g/mol (B) 52 g/mol (C) 70 g/mol (D) 86 g/mol
17. The vapor pressure of pure water at 60 °C is 149 torr. What is the approximate vapor pressure of a solution prepared by adding 135 grams of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) to 300 grams of water at 60 °C?
- (A) 137.4 torr (B) 102.8 torr (C) 142.6 torr (D) 6.5 torr
18. Which one of the following solutions has the **lowest** boiling point?
- (A) 0.10 m $\text{Mg}_3(\text{PO}_4)_2$ (B) 0.32 m NH_4Cl
(C) 0.20 m Na_2SO_4 (D) 0.11 m K_3AsO_4
19. Consider the strong electrolytes, sodium sulfate, Na_2SO_4 , and sodium phosphate, Na_3PO_4 .
- If 350 mL of 0.30 M $\text{Na}_2\text{SO}_4(\text{aq})$ is added to 250 mL of 0.50 M $\text{Na}_3\text{PO}_4(\text{aq})$, the sodium ion concentration (i.e Molarity) in the mixture, $[\text{Na}^+]$, is approximately:
- (A) 0.59 M (B) 0.046 M (C) 0.98 M (D) 0.35 M

PART II. TWO (2) PROBLEMS ON FOLLOWING PAGES:

REMEMBER TO SHOW YOUR WORK FOR CREDIT

Version B

- (12) 1. Consider the equilibrium between $\text{N}_2(\text{g})$, $\text{H}_2(\text{g})$, $\text{NH}_3(\text{g})$:
 $\text{N}_2(\text{g}) + 3 \text{H}_2(\text{g}) \rightleftharpoons 2 \text{NH}_3(\text{g})$. The value of K_c at $30\text{ }^\circ\text{C}$ is 8.0. The Enthalpy change for this reaction is $\Delta H = -92.0\text{ kJ}$.

Calculate the temperature, **in $^\circ\text{C}$** , at which the equilibrium constant, K_c , is 1.00×10^{-4} .

- (12) 2. The vapor pressure of pure Ethylbenzene, $\text{C}_6\text{H}_5\text{C}_2\text{H}_5(\text{l})$ [$M=106$], is 74.0 torr at $70\text{ }^\circ\text{C}$. When 50 grams of an unknown non-volatile solute, X, is added to 265 grams of Ethylbenzene, the vapor pressure of the solution at $70\text{ }^\circ\text{C}$ is 62.9 torr.

Calculate the Molar Mass of the unknown, X, in grams/mol

