

CHEM 1423 - Exam 3 – March 30, 2017 - Version A

Name _____

(76) PART I. MULTIPLE CHOICE (Circle the ONE correct answer)

1. What is the approximate pH of a solution containing 0.07 grams of the strong base, potassium hydroxide, KOH ($M = 56.1 \text{ g/mol}$), dissolved in 50 L of aqueous solution?
(A) 4.3 (B) 4.6 (C) 9.7 (D) 9.4
2. Which of the following aqueous solutions is/are acidic?
(i) Ammonium Bromide (NH_4Br)
(ii) Potassium Bromide (KBr)
(iii) Anilinium Chloride (AnilHCl)
(iv) Sodium Lactate (NaLac)
(A) iv only (B) i & iii (C) i & ii & iv (D) i & ii & iii
3. What is the Base Equilibrium Constant of the weak base, Hydroxylamine, if a 0.05 M hydroxylamine solution has a $\text{pH} = 9.4$?
(A) 1.3×10^{-8} (B) 3.2×10^{-18} (C) 5.0×10^{-4} (D) 4.8×10^{-6}

For #4 - #5: Consider the weak base, Morphine (Morp). Its base equilibrium constant is 1.6×10^{-6} .

4. What is the approximate **percent protonation** in a 0.07 M solution of Morphine?
(A) $3.3 \times 10^{-2} \%$ (B) $4.8 \times 10^{-3} \%$ (C) $3.3 \times 10^{-4} \%$ (D) 0.48%
5. What is the approximate pH of a 0.04 M solution of Morphinium Chloride, (MorpHCl) ?
(A) 4.8 (B) 3.6 (C) 5.3 (D) 9.2
6. Benzoic Acid (HBenz) is a weak acid with an acid dissociation constant of 6.3×10^{-5} . What is the approximate pH of a 0.20 M solution of Sodium Benzoate (NaBenz)?
(A) 2.4 (B) 5.2 (C) 8.8 (D) 11.6
7. Sulfurous Acid, H_2SO_3 , is a diprotic acid with acid dissociation constants, $K_{a1} = 1.2 \times 10^{-2}$, and $K_{a2} = 6.6 \times 10^{-8}$. What is the approximate pH of a 0.10 M solution of sodium sulfite, Na_2SO_3 ?
(A) 3.9 (B) 10.1 (C) 9.9 (D) 4.1

Version A

8. Arsenous Acid, HAsO_2 , has an acid dissociation constant, $K_a = 8.0 \times 10^{-10}$. What is the approximate percent dissociation of a 0.03 M solution of HAsO_2 ?
- (A) $4.9 \times 10^{-4} \%$ (B) $1.6 \times 10^{-2} \%$ (C) 9.9 (D) $1.6 \times 10^{-4} \%$
9. If added to 2 L of 0.50 M NaOH, which one of the following would form a buffer?
- (A) 2. L of 0.40 M Nitric Acid (HNO_3)
(B) 2. L of 0.40 M Potassium Acetate (KAc)
(C) 2. L of 0.60 M Lactic Acid (HLac)
(D) 2. L of 0.40 M Acetic Acid (HAc)
10. The base equilibrium constant for aniline (Anil) is 4.3×10^{-10} . What is the approximate pH of a solution containing 0.30 M Aniline and 0.50 M Anilinium Bromide (AnilHBr)?
- (A) 9.60 (B) 4.85 (C) 4.41 (D) 9.15
11. The acid dissociation constant of Propanoic Acid (HProp) is 1.5×10^{-5} . What is the approximate pH after 0.40 mol of NaOH is added to a solution initially containing 2.0 L of 0.50 M HProp ?
- (A) 4.65 (B) 5.00 (C) 4.42 (D) 5.22

For #12 - #14: Consider the amino acid, Glutamic Acid (Glu). The most positive form of Glutamic Acid is Glu^{1+} and the most negative form is Glu^{2-} . The three pK_a 's of Glutamic Acid are: $\text{pK}_a' = 2.1$, $\text{pK}_a'' = 4.1$, and $\text{pK}_a''' = 9.5$.

12. What is the isoelectric point (pI) of Glutamic Acid?
- (A) 3.1 (B) 6.8 (C) 2.1 (D) 4.1
13. At what pH does one have 50% Glu^0 (neutral) and 50% Glu^{1-} ?
- (A) 3.1 (B) 9.5 (C) 6.8 (D) 4.1
14. What is the average charge on the Glutamic Acid molecule at $\text{pH} = 6.8$?
- (A) -1.5 (B) -0.5 (C) -1.0 (D) -2.0

Version A

15. 240 mL of 0.40 M NaOH(aq) is needed to completely neutralize 120 mL of an aqueous $\text{H}_3\text{PO}_4(\text{aq})$ solution? What is the Molarity of the $\text{H}_3\text{PO}_4(\text{aq})$ solution?
- (A) 0.27 M (B) 0.47 M (C) 0.80 M (D) 0.21 M

For #16 - #19: Arsenous acid, H_3AsO_3 , is a triprotic acid with acid dissociation constants, $K_a' = 5.6 \times 10^{-3}$, $K_a'' = 1.7 \times 10^{-7}$, and $K_a''' = 2.5 \times 10^{-12}$

16. Which one of the solutions below would be best to prepare a buffer with $\text{pH} = 7.2$?
- (A) $\text{K}_2\text{HAsO}_3/\text{Na}_3\text{AsO}_3$ (B) Na_2HAsO_3
(C) $\text{KH}_2\text{AsO}_3/\text{Na}_2\text{HAsO}_3$ (D) $\text{KH}_2\text{AsO}_3/\text{Na}_3\text{AsO}_3$
17. What is the approximate pH of a solution containing pure K_2HAsO_3 ?
- (A) 6.8 (B) 9.2 (C) 11.6
(D) The pH depends upon the K_2HAsO_3 concentration.
18. What is the approximate pH of a solution prepared by adding 0.30 mol of NaOH to 2.0 L of 0.40 M KH_2AsO_3 ?
- (A) 7.20 (B) 7.00 (C) 6.34 (D) 6.55
19. Approximately what ratio of $[\text{H}_3\text{AsO}_3]/[\text{H}_2\text{AsO}_3^-]$ will give a solution with $\text{pH} = 2.9$?
- (A) 0.2 (B) 0.6 (C) 5.6 (D) 4.5

PART II. TWO (2) PROBLEMS BELOW: REMEMBER TO SHOW WORK FOR CREDIT

Versions A and B

- (16) 1. Tellurous acid, H_2TeO_3 , is a diprotic acid with acid dissociation constants, $K_{a'} = 3.0 \times 10^{-3}$ and $K_{a''} = 2.0 \times 10^{-8}$
- (6) (a) Calculate the pH of a solution prepared by mixing 2.0 L of 0.60 M K_2TeO_3 with 1.0 L of 0.70 M HCl.
- (6) (b) Calculate the pH of a solution prepared by mixing 2.0 L of 0.60 M H_2TeO_3 with 2.0 L of 0.80 M NaOH.
- (4) (c) Calculate the ratio, $[\text{HTeO}_3^-]/[\text{H}_2\text{TeO}_3]$, needed to prepare a buffer solution with $\text{pH} = 3.20$
- (08) 2. Hypobromous Acid, HBrO , has an Acid Dissociation Constant, $K_a = 2.0 \times 10^{-9}$. Calculate the pH of a solution of Sodium Hypobromite, NaBrO ($M = 118.9 \text{ g/mol}$) that is 3.0% NaBrO by weight.
- Note:** Assume that the density of the solution is 1.0 g/mL

Version B

For #15 - #18: Arsenous acid, H_3AsO_3 , is a triprotic acid with acid dissociation constants, $K_a' = 5.6 \times 10^{-3}$, $K_a'' = 1.7 \times 10^{-7}$, and $K_a''' = 2.5 \times 10^{-12}$

15. What is the approximate pH of a solution containing pure K_2HAsO_3 ?
(A) 9.2 (B) 11.6 (C) 6.8
(D) The pH depends upon the K_2HAsO_3 concentration.
16. Which one of the solutions below would be best to prepare a buffer with $\text{pH} = 7.2$?
(A) $\text{K}_2\text{HAsO}_3/\text{Na}_3\text{AsO}_3$ (B) Na_2HAsO_3
(C) $\text{KH}_2\text{AsO}_3/\text{Na}_3\text{AsO}_3$ (D) $\text{KH}_2\text{AsO}_3/\text{Na}_2\text{HAsO}_3$
17. What is the approximate pH of a solution prepared by adding 0.30 mol of NaOH to 2.0 L of 0.40 M KH_2AsO_3 ?
(A) 6.55 (B) 7.00 (C) 6.34 (D) 7.20
18. Approximately what ratio of $[\text{H}_3\text{AsO}_3]/[\text{H}_2\text{AsO}_3^-]$ will give a solution with $\text{pH} = 2.9$?
(A) 2.5 (B) 0.6 (C) 5.6 (D) 0.2
19. 240 mL of 0.40 M NaOH(aq) is needed to completely neutralize 120 mL of an aqueous H_3PO_4 (aq) solution? What is the Molarity of the H_3PO_4 (aq) solution?
(A) 0.47 M (B) 0.80 M (C) 0.27 M (D) 0.21 M

PART II. TWO (2) PROBLEMS BELOW: REMEMBER TO SHOW WORK FOR CREDIT

Versions A and B

- (16) 1. Tellurous acid, H_2TeO_3 , is a diprotic acid with acid dissociation constants, $K_{a'} = 3.0 \times 10^{-3}$ and $K_{a''} = 2.0 \times 10^{-8}$
- (6) (a) Calculate the pH of a solution prepared by mixing 2.0 L of 0.60 M K_2TeO_3 with 1.0 L of 0.70 M HCl.
- (6) (b) Calculate the pH of a solution prepared by mixing 2.0 L of 0.60 M H_2TeO_3 with 2.0 L of 0.80 M NaOH.
- (4) (c) Calculate the ratio, $[\text{HTeO}_3^-]/[\text{H}_2\text{TeO}_3]$, needed to prepare a buffer solution with $\text{pH} = 3.20$
- (08) 2. Hypobromous Acid, HBrO , has an Acid Dissociation Constant, $K_a = 2.0 \times 10^{-9}$. Calculate the pH of a solution of Sodium Hypobromite, NaBrO ($M = 118.9 \text{ g/mol}$) that is 3.0% NaBrO by weight.

Note: Assume that the density of the solution is 1.0 g/mL