

CHEM 1423
Chapter 19
Homework Questions

TEXTBOOK HOMEWORK

19.12 Find the pH of a buffer that consists of 1.3 M sodium phenolate ($\text{C}_6\text{H}_5\text{ONa}$) and 1.2 M phenol ($\text{C}_6\text{H}_5\text{OH}$) (pK_a of phenol = 10.00).

19.15 Find the pH of a buffer that consists of 0.50 M methylamine (CH_3NH_2) and 0.60 M $\text{CH}_3\text{NH}_3\text{Cl}$ (pK_b of CH_3NH_2 = 3.35).

19.16 What is the component concentration ratio, $[\text{Pr}^-]/[\text{HPr}]$, of a buffer that has a pH of 5.44 (K_a of HPr = 1.3×10^{-5})?

19.18 A buffer containing 0.2000 M of acid, HA, and 0.1500 M of its conjugate base, A^- , has a pH of 3.35. What is the pH after 0.0015 mol of NaOH is added to 0.5000 L of this solution?

19.19 A buffer that contains 0.40 M base, B, and 0.25 M of its conjugate acid, BH^+ , has a pH of 8.88. What is the pH after 0.0020 mol of HCl is added to 0.25 L of this solution?

19.42(C) Find the pH during the titration of 20.00 mL of 0.1000 M triethylamine, $(\text{CH}_3\text{CH}_2)_3\text{N}$ (K_b = 5.2×10^{-4}), with 0.1000 M HCl solution after the addition of 15.00 mL of titrant.

19.50 The solubility of silver carbonate is 0.032 M at 20 °C. Calculate its K_{sp} .

19.56 Calculate the molar solubility of $\text{Ca}(\text{IO}_3)_2$ in (a) 0.060 M $\text{Ca}(\text{NO}_3)_2$ and (b) 0.060 M NaIO_3 . (See Appendix C.). Also calculate the molar solubility in pure water.

SUPPLEMENTARY HOMEWORK

Buffers

S1. In a buffer solution, if $[A^-] > [HA]$, which of the following must be true?

- a. $pH < pK_a$
- b. $pH = pK_a$
- c. $pH > pK_a$
- d. $pH < 7.00$
- e. $pH > 7.00$

S2. A buffer solution is 0.080 M in lactic acid ($K_a = 1.8 \times 10^{-4}$) and 0.070 M in sodium lactate. The pH of the solution is

- a. 2.86.
- b. 3.68.
- c. 3.80.
- d. 4.18.
- e. 4.62.

S3. Which acid, in combination with its conjugate base, would be the best choice to make a buffer of $pH = 4.35$?

- a. acetic acid ($K_a = 1.8 \times 10^{-5}$)
- b. benzoic acid ($K_a = 6.3 \times 10^{-5}$)
- c. formic acid ($K_a = 1.8 \times 10^{-4}$)
- d. hydrofluoric ($K_a = 7.2 \times 10^{-4}$)
- e. nitrous acid ($K_a = 4.5 \times 10^{-4}$)

S4. If a buffer is made up using 1.00 mole of a weak acid ($pK_a = 5.00$) and 0.90 mole of its conjugate base, which of the following must be true?

- a. $pH < 5.00$
- b. $pH = 5.00$
- c. $pH > 5.00$
- d. $pH = 7.00$
- e. $pH > 7.00$

S5. For each of the solutions below, indicate whether the solution would be a buffer.

- a) A solution prepared by adding 1 L of 0.50 M NaOH to 2 L of 0.5 M HAc (acetic acid)

- b) A solution prepared by adding 1 L of 1.0 M HCl to 2 L of 0.80 M NaAc (sodium acetate)

- c) A solution prepared by adding 1 L of 0.50 M H_2CO_3 to 1 L of 0.50 M K_2CO_3

- d) A solution prepared by adding 1 L of 0.50 M HCl to 2 L of 0.50 M K_2CO_3

- e) A solution prepared by adding 3 L of 0.50 M HCl to 2 L of 0.50 M K_2CO_3

- f) A solution prepared by adding 5 L of 0.50 N HCl to 2 L of 0.50 M K_2CO_3

- g) A solution prepared by adding 1 L of 0.25 M HNO_3 to 1 L of 0.50 M K_2Prop (potassium propanoate)

- h) A solution prepared by adding 1 L of 0.25 M NaOH to 1 L of 0.50 M K_2Prop (potassium propanoate)

S6. The following **independent** questions are on pH calculations in solutions of Arsenic Acid (H_3AsO_4) and its various anions. H_3AsO_4 is a triprotic acid with Acid Dissociation Constants: $K_a' = 6.0 \times 10^{-3}$, $K_a'' = 1.0 \times 10^{-7}$, $K_a''' = 3.2 \times 10^{-12}$

- a) Calculate the pH of a solution prepared by mixing 3.0 L of 0.40 M H_3AsO_4 with 1.0 L of 0.80 M KOH.
- b) Calculate the pH of a solution prepared by mixing 3.0 L of 0.40 M H_3AsO_4 with 2.0 L of 0.80 M KOH.
- c) Calculate the pH of a solution prepared by mixing 3.0 L of 0.40 M H_3AsO_4 with 4.0 L of 0.80 M KOH.
- d) Calculate the pH of a solution prepared by mixing 3.0 L of 0.40 M Na_3AsO_4 with 1.0 L of 0.80 M HNO_3 .
- e) Calculate the pH of a solution prepared by mixing 3.0 L of 0.40 M Na_3AsO_4 with 2.0 L of 0.80 M HNO_3 .
- f) Calculate the pH of a solution prepared by mixing 3.0 L of 0.40 M Na_3AsO_4 with 4.0 L of 0.80 M HNO_3 .
- g) What will be the pH of a buffer solution containing $[\text{H}_2\text{AsO}_4^-] = 0.40$ M and $[\text{HAsO}_4^{2-}] = 0.60$ M?
- h) What will be the pH of a buffer solution containing $[\text{HAsO}_4^{2-}] = 0.40$ M and $[\text{AsO}_4^{3-}] = 0.15$ M?
- i) What value of the ratio, $[\text{H}_2\text{AsO}_4^-]/[\text{H}_3\text{AsO}_4]$, is required to prepare a buffer with $\text{pH} = 2.60$?
- j) What value of the ratio, $[\text{HAsO}_4^{2-}]/[\text{AsO}_4^{3-}]$, is required to prepare a buffer with $\text{pH} = 10.90$?

Titration

S7. 138. mL of 0.105 M KOH was required to completely titrate a 25.00 mL sample of H_3PO_4 .

Determine the Molarity of the H_3PO_4 sample.

S8. Consider the weak base, pyridine [$\text{C}_5\text{H}_5\text{N} \equiv \text{Pyr}$], which has $K_b = 1.7 \times 10^{-9}$. Titration of pyridine with a strong acid causes formation of the pyridinium ion [$\text{C}_5\text{H}_5\text{NH}^+ \equiv \text{PyrH}^+$].

(a) Calculate the pH of a solution formed by adding 30.00 mL of 0.20 M HNO_3 to 50.00 mL of 0.20 M pyridine.

(b) Calculate the pH of a solution formed by adding 50.00 mL of 0.20 M HNO_3 to 50.00 mL of 0.20 M pyridine.

S9. Vitamin C ($\text{C}_6\text{H}_8\text{O}_6$, $M = 176$.) is a monoprotic acid. To analyze a Vitamin C capsule weighing 0.64 grams by titration, 23.6 mL of 0.120 M NaOH was required. Calculate the mass percent of Vitamin C in the capsule.

S10. When 6.00 grams of a sample of impure Strontium Hydroxide [$\text{Sr}(\text{OH})_2$, $M = 121.6$] is titrated with 0.340 M HNO_3 , it takes 224. mL of the strong acid to completely titrate the base. Calculate the mass percent of impurity in the Strontium Hydroxide sample.

Solubility Product

S11. The solubility product of the slightly soluble salt, Ag_2CO_3 , is $K_{sp} = 6.2 \times 10^{-12}$.

a) What is the solubility and the $[\text{Ag}^+]$ and $[\text{CO}_3^{2-}]$ concentrations in pure water?

b) What is the solubility in a solution containing Ag_2CO_3 and 0.20 M AgNO_3 ?

c) What is the concentration of silver ions, $[\text{Ag}^+]$ in a solution containing Ag_2CO_3 and 0.10 M K_2CO_3 ?

S12. The solubility products of two sparingly soluble Bromide (Br^-) salts are:




Consider a solution which initially contains $5.0 \times 10^{-5} \text{ M Ag}^+(\text{aq})$ and $5.0 \times 10^{-5} \text{ M Hg}^{2+}(\text{aq})$. KBr (a strong electrolyte) is added until $[\text{Br}^-] = 2.0 \times 10^{-8} \text{ M}$. Which of the above salts (AgBr and HgBr_2) will precipitate?

S13. Mercury(I) Sulfate, Hg_2SO_4 , is a sparingly soluble salt with $K_{\text{sp}} = 6.5 \times 10^{-7}$.


If 1200 mL of 0.010 M $\text{K}_2\text{SO}_4(\text{aq})$ is mixed with 800 mL of 0.020 M $\text{HgNO}_3(\text{aq})$, calculate the concentrations of $[\text{Hg}^+]$ and $[\text{SO}_4^{2-}]$ in the resulting solution and **determine whether or not** $\text{Hg}_2\text{SO}_4(\text{s})$ will precipitate.

Amino Acids

S14. Consider the amino acid, Lysine. The most positive and most negative forms are shown below:



Lys²⁺



Lys⁻

The three pK_a 's are: $\text{pK}_a'(\alpha\text{-COOH}) = 2.18$ $\text{pK}_a''(\alpha\text{-NH}_3^+) = 8.95$ $\text{pK}_a'''(\epsilon\text{-NH}_3^+) = 10.53$

- What is the isoelectric point?
- At what pH does one have 100% Lys^+ .
- At what pH does one have 50% Lys and 50% Lys^-
- What is the composition of the solution at $\text{pH} = 2.18$

